

Ex. № 11
від **11 .10.2024р.**

Medical Reference
given to: Protsenko Oleksandr

After the clinical examination at the clinic, she received a prescription for the treatment of the existing disease.

Diagnosis:

Dyscirculatory encephalopathy of the II degree of mixed vascular and post-traumatic genesis (traumatic brain injury in 2015) in the form of astheno-neurotic, hypertensive, cephalic, tinnitus, vestibulo-cerebellar syndromes. Diabetes mellitus of type 2. Arterial hypertension of the 2nd degree. Urinary stone disease.

It is recommended to take individually selected angiotherapeutic mixtures in order to continue treatment in an outpatient setting for an optimal sanogenic effect.

Medicinal mixtures are given: 1 syringe of 8 ml for taking it oral, 15 syringes of 2 ml for intramuscular injection in the morning according to the agreed schedule, 15 syringes of 3 ml for intramuscular injection in the evening. Be sure to store in the refrigerator. The storage conditions for the transportation of medicinal mixtures by any means of transport are: refrigerator or in a refrigerator bag with ice cartridges. The container of medicinal mixtures is in syringes. Do not change the container during transportation and further use in order to avoid oxidation and chemical reaction, which will lead to the deterioration of medicinal mixtures.

Medical Director



U.B. Lushchyk

